



COMPARATIVE EVALUATION OF GEL BAITS PALATABILITY AGAINST *BLATTELLA GERMANICA*

¹MARCOS R. POTENZA, ¹FABRÍCIO C. REIS, ¹THAIS L. LIMA; ²WASHINGTON L. B. FERREIRA AND ²GUILHERME STAGNI

¹Av. Cons. Rodrigues Alves, 1252, São Paulo/SP, 04014 002, Brazil/ Instituto Biológico/APTA/SAA
E-mail: potenza@biologico.sp.gov.br

²Av. Nações Unidas, 18001 4º andar São Paulo/SP 04795 900, Brazil/ Syngenta, Professional Pest Management LATAM



INTRODUCTION

The control of *Blattella germanica* represents a major challenge for Brazilian PCOs (Pest Control Operators), due to their great capacity for reproduction and dispersal on tropical weather environments. Several gel baits insecticides have been developed and sold in Brazil, during the past few years, from a wide amount of multinational and local manufacturers. Although the active ingredient is a key component on a bait, the food matrix is also a fundamental part of a product and it may vary from manufacturer to manufacturer due to the alimentary components used. The quality of the bait food matrix is directly related to its performance on the most challenging situations, such as kitchens and food processing areas, especially when competing with other food sources.

OBJECTIVES

The aim of this study was to compare the palatability of 9 distinct gel baits against a new product on Brazilian market – Advion Cockroach Gel Bait® (Indoxacarb 0.6%) – when offered to a population of *Blattella germanica*.

MATERIAL AND METHODS

The products tested against Advion®, named here as P1, were: P2: a Hydramethylnon (2%) gel bait, P3: a Hydramethylnon (2%) gel bait, P4: a Fipronil (0.05%) gel bait, P5: a Fipronil (0.05%) gel bait, P6: a Imidacloprid (2.15%) gel bait, P7: a Imidacloprid (2.15%) gel bait, P8: a Imidacloprid (2.15%) gel bait, P9: a Imidacloprid (2.15%) gel bait and P10: a Indoxacarb (0.6%) gel bait, and all of them were present at Brazilian market by the time the trials were made. Ten repetitions were made for each P1 x Other Product trial. In an arena (33cm length, 22cm width, 10cm height) containing 30 individuals of *Blattella germanica* – 10 male, 10 female and 10 nymphs, fasted for 24 hours before the trials – 0.5g of P1 and 0.5g of other gel bait were offered simultaneously for 2 hours (Figure 1). After this period, both remaining baits were weighted on a high precision balance.

RESULTS

Results showed that P1 has superior palatability when compared to all its competitors, as shown on Table 1. The obtained information is valuable when designing an effective cockroach control program where a palatable gel bait formulation is required.

Table 1. Comparative evaluation of gel baits palatability against *Blattella germanica*.




Fig. 1. Two gel baits being offered simultaneously to *Blattella germanica*.



Published in ICUP 2017 Proceedings, available from QR code & www.icup.org.uk

