

TROPICAL HOUSEHOLD ANTS: PEST STATUS, SPECIES DIVERSITY, FORAGING BEHAVIOR, and BAITING STUDIES

Chow-Yang Lee

Urban Entomology Laboratory, Vector Control Research Unit, School of Biological Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia

Abstract Questionnaire-based surveys of homeowners revealed that ants were the most economically important and abundant household pest after mosquitoes and cockroaches in Malaysia. Twenty-five species of ants were found indoors and outside buildings; all species nest outdoors except *Monomorium pharaonis*, *Monomorium floricola*, *Tapinoma melanocephalum*, and *Solenopsis molesta*. Surveys of houses and food stores indicated the most common species were *Pheidole* sp., *T. melanocephalum*, *Monomorium destructor*, and *Paratrechina longicornis*. A survey of homeowners' attitudes and knowledge of ants showed that 62% found ants daily in their homes. Most respondents tolerated < 50 ants indoors; 65% used aerosol sprays for control, 6% used ant baits. Homeowners were not aware that ants can mechanically vector pathogenic organisms. Bacteria, fungi, and yeasts were isolated from ants collected from food outlets. Most ant species responded to peanut butter attractant, except *P. longicornis* and *T. melanocephalum*, which preferred honey. Attractancy to peanut butter and honey changed in an 18-month study of *M. pharaonis*. All species, except *Pheidole* sp. preferred liquid bait. Foraging of *P. longicornis*, *M. pharaonis*, and *Solenopsis geminata* was negatively correlated with temperature; their peak activity was 2-4 hours after sunset. Most baits gave >75% reduction after one week post-treatment.

Key Words household ants, Malaysia, pest status, food preference, foraging, baits

INTRODUCTION

Household ants are an important group of insect pests in the urban environment because of their close association with mankind (Holl Dobler and Wilson, 1990) and foraging activity in mass numbers (Chong, 1997; Robinson, 1996; Lee et al., 1999). Of the 10,000 ant species that have been described, less than 0.5% are pests in the human environment, particularly in structures and buildings in south Asia (Lee and Robinson, 2001). Prior to the 1990s, household ants were considered a less important group of household pests than cockroaches and mosquitoes in Asia, but their status has risen in many Asian developed countries, such as South Korea and Singapore. In tropical Malaysia, ant control accounted for about 10% of the business of the pest control industry in 1995 (Yap and Lee, 1996), while in other developed countries in Asia, it has a market share of between 15-30%.

The species of household ants that have been studied extensively include Pharaoh ant, *Monomorium pharaonis*, (Edwards, 1986), carpenter ants, *Camponotus* spp., (Hansen and Akre, 1993), Argentine ant, *Linepithema humile*, (Vega and Rust, 2001), and fire ant, *Solenopsis* spp. (Taber, 2000; Williams et al., 2001). Information is limited on important biological aspects and behaviour of many tropical species of household ants such as ghost ant, *Tapinoma melanocephalum*, crazy ant, *Paratrechina longicornis*, and big-headed ant, *Pheidole megacephala*. The objective of this paper is to summarize some of the published and unpublished findings of research projects on tropical household ants conducted in Malaysia. Presented here will be pest status and human perception, and some important biological aspects, such as species composition, distribution, seasonal food preference, and foraging patterns of household ants in the tropics.

Household Pest Status

Several questionnaire-based surveys on the status of household pests were done between 1983-2001 among homeowners and the pest control industry in Malaysia (Table 1). The survey conducted in 1983 (Yap and Foo, 1984) revealed that ants were the most important group of household pests after mosquitoes and cockroaches. A similar finding was obtained in 1995. In the survey conducted in 1998 (Yap et al., 1999) ants were still at the same position as before, but cockroaches have become the most important household pests, rather than mosquitoes. Currently, improved household insecticide products are available for mosquitoes compared to products for other household insect pests in Malaysia. Products for cockroach and ant control are primarily limited to aerosols, and only three cockroach baits and one ant bait are available. Estimated figures show that Malaysian households spent US\$48 million annually on household insecticide products in 1997/1998 (Yap, 1999) and US\$57 million in 2000/2001 (I. Ridzuan, personal communication), compared to about US\$25 million in 1986/1987 (Yap, 1988). Only 5-10% of these figures were spent on products for cockroach and ant control.

A survey in 2001 of the importance of pest groups to the Malaysian pest control industry showed that household ant control ranked third after termite and cockroach control (C.Y. Lee, unpublished), compared to its fourth position in a 1995 survey (Yap and Lee, 1996) (Table 2). Currently, it accounts for 10% of an estimated total business volume of US\$18-20 million (Lee et al., 1999). The demand for household ant control in Malaysia, especially in food preparative outlets has increased since the introduction of cockroach gel baits for German cockroach control in 1997. In many countries around the world, including the United States, ants have become the number one household pest in terms of control revenue (Gooch, 1999; Kaminski, 2000; Jenkins, 2001), and rated the most difficult pests to control (Gooch, 1999).

Structure-Invading Ants

Yap and Lee (1994) reported on household ants in residential premises in Penang, Malaysia. They recorded *M. pharaonis* as the predominant species, followed by *Tapinoma sessile* and *P. longicornis*. Between June 2000 and December 2001, 479 households were sampled in Penang, Malaysia; these sites ranged from houses to apartments in rural, suburban, and urban areas. Ants were collected from baited cards (6.2 by 7.5 cm) placed in the living room, dining area and kitchen, along the perimeter or corridor of the house, and outside the house. Twenty-five ant species from four subfamilies – Myrmicinae, Formicinae, Dolichoderinae, and Ponerinae – were collected (Table 3). With the exception of *M. pharaonis*, *M. floricola*, *T. melanocephalum*, and *S. molesta*,

Table 1. Household pest status in Penang, Malaysia (1983 – 1998)

Household pest	Year of survey		
	1983 ¹	1995 ²	1998 ³
Cockroaches	2	2	1
Ants	3	3	3
Mosquitoes	1	1	2
Rodents	4	4	4
House flies	5	5	5
Others ⁴	6	6	6

¹Based on Yap and Foo (1984) (n = 395 houses).

²H.H. Yap and C.Y. Lee, unpublished data (n = 814 houses).

³Yap et al. (1999) (n = 245 houses).

⁴Others = geckos, spiders, wasps, fleas, begbugs, etc.

Table 2. Rank of targeted pests of the pest control industry in Malaysia (based on total business volume)

Pest	Year of survey	
	1995 ¹	2001 ²
Termites	1	1
Cockroaches	2	2
Ants	4	3
Rodents	3	4
Mosquitoes	4	5
Flies & other incidental pests	6	6

¹Yap and Lee (1996) (n = 42 companies)

²C.Y. Lee, unpublished data. (n = 55 companies).

Table 3. Household ants of Malaysia – locations of collection (Na & Lee, 2001)

Subfamily Species	Common name	Location where ants were found within a structure/building		
		Indoors	Perimeter	Outdoors
Subfamily Myrmicinae				
<i>Monomorium pharaonis</i>	Pharaoh ant	x	x	-
<i>Monomorium destructor</i>	Singapore ant	x	x	-
<i>Monomorium orientale</i>	—	x	x	-
<i>Monomorium minimum</i>	—	x	x	-
<i>Monomorium floricola</i>	—	-	x	-
<i>Crematogaster</i> sp.	Acrobat ant	-	x	-
<i>Pheidole</i> sp.	Big-headed ant	x	-	-
<i>Tetramorium</i> sp.	Pavement ant	-	x	-
<i>Solenopsis geminata</i>	Tropical fire ant	x	x	x
<i>Solenopsis invicta</i>	Imported fire ant	x	x	x
<i>Solenopsis molesta</i>	Thief ant	x	-	-
Subfamily Formicinae				
<i>Anoplolepis longipes</i>	Red crazy ant	-	x	-
<i>Paratrechina longicornis</i>	Crazy ant	x	x	x
<i>Paratrechina</i> sp.	Crazy ant	-	x	x
<i>Camponotus</i> sp.	Carpenter ant	-	x	x
<i>Prenolepis imparis</i>	Small honey ant	-	x	-
<i>Formica</i> sp.	Field ant	-	x	x
Subfamily Dolichoderinae				
<i>Tapinoma sessile</i>	Odorous house ant	x	x	x
<i>Tapinoma melanocephalum</i>	Ghost ant	x	x	-
<i>Tapinoma indicum</i>	Ghost ant	x	x	-
<i>Linepithema humile</i>	Argentine ant	x	x	-
<i>Oecophylla smaragdina</i>	Weaver ant	-	-	x
<i>Technomyrmex albipes</i>	White-footed ant	-	x	-
<i>Dolichoderus bituberculata</i>	Rambutan ant	x	x	x
Subfamily Ponerinae				
<i>Odontoponera</i> sp.	—	-	x	x

which nest indoors, the majority of the ants collected were outdoor-nesting species, but forage indoors for food.

The most dominant species trapped in the residential premises was the big-headed ant, *Pheidole* sp., followed by the ghost ant, *T. melanocephalum*, the Singapore ant, *Monomorium destructor*, and the crazy ant, *P. longicornis* (Table 4). The big-headed ant, which is also an important insect pest in agriculture (Reimer et al., 1990), was often found nesting outdoors under flower pots. The ghost ant primarily occurs outdoors, but many nests were reported in kitchens. Nesting habitats for *M. destructor* are generally outdoors, especially in soil with vegetation and shrubs, while *M. pharaonis* and *M. floricola* prefer to nest indoors.

Surveys were conducted on ants in food outlets, including university cafeterias, coffee shops, residential kitchens, and hotel kitchens (Lee et al. 2001a), and university dormitory pantries (Loke, 2002). In both studies, *M. destructor* was the dominant species (Table 4). Other species found in large numbers were *P. longicornis*, *Pheidole* sp., and *T. melanocephalum*. The university cafeterias showed the highest diversity of ant species; species trapped were: *P. longicornis*, *T. melanocephalum*, and *Monomorium* spp. They are cosmopolitan in distribution and considered

Table 4. Survey on household ants in residential premises and food-handling outlets

Species	% of total ants collected		
	Residential ¹ (n = 479)	Food outlets ² (n = 31)	Dormitory pantries ³ (n = 124)
<i>Anoplolepis longipes</i>	0.4	1.7	6.8
<i>Camponotus</i> sp.	<0.1	0.1	<0.1
<i>Dolichoderus bituberculata</i>	0.1	1.2	—
<i>Formica</i>	—	—	0.2
<i>Monomorium destructor</i>	9.1	27.8	25.1
<i>Monomorium floricola</i>	3.0	10.8	0.6
<i>Monomorium minimum</i>	2.0	—	—
<i>Monomorium pharaonis</i>	6.2	3.4	7.4
<i>Odontoponera</i> sp.	<0.1	—	—
<i>Paratrechina longicornis</i>	4.9	9.0	21.2
<i>Pheidole</i> sp.	32.4	16.4	21.7
<i>Prenolepis imparis</i>	0.2	0.3	0.9
<i>Solenopsis geminata</i>	0.1	3.7	—
<i>Solenopsis invicta</i>	1.9	—	3.6
<i>Tapinoma melanocephalum</i>	27.7	19.5	12.3
<i>Tapinoma indicum</i>	8.9	4.2	0.2
<i>Tetramorium</i> sp.	0.1	1.9	—
Others	2.9	—	—

¹Based on Na (2001) and Lee, unpublished data (n = 52,389 ants).

²Based on Lee et al. (2001) (n = 5531 ants).

³Based on Loke (2002) (n = 19,083 ants).

to be tramp ants. Tramp ants are polygynous, unicolonial, reproduce by budding, largely dispersed worldwide through human commerce, and live in close association with humans (Holldobler and Wilson, 1990; Passera, 1994). Although a number of ant species demonstrate anthropophilic tendencies (Pisarski, 1982; Prins et al., 1990; Thompson, 1990), only tramp species are truly domestic with great ability for nest changes. Nesting in an unstable habitat such as the human environment requires frequent migration, and migration has become a unique characteristic of tramp ant species (Passera, 1994). The crazy ant, *P. longicornis*, is often observed to be the first species to succeed in a disturbed habitat or a new building or structure (C.Y. Lee, unpublished). Tramp ant species often displace other ant species once they become established. Wetterer et al. (1999) reported the displacement of several species and other insects by *P. longicornis* in a greenhouse structure.

Homeowner Attitude and Knowledge Concerning Household Ants

Wood et al. (1981) reported that an understanding of the attitude and knowledge of homeowners toward pests is important to the success of a pest management program, especially in terms of their appreciation of the value of sanitation and pesticide application. To have a better understanding of the perception of the Malaysia homeowners towards ant problems, a questionnaire-based survey was conducted in Penang, Malaysia. Penang has a population of about 1 million and is located in northern Malaysia. Face-to-face interviews with 256 residents of single-story, double-story, link houses, and apartments in urban, suburban, and rural areas were conducted between May and September 2000.

A total of 159 homeowners reported ants and ant problems in their houses (Table 5); there was no significant difference between ($\chi^2 = 2.44$, $P > 0.05$, $df = 3$) between these two responses. When asked about the cause(s) of ants in their home, 70% reported food and beverages as the

Table 5. Survey on attitude and knowledge of homeowners toward ant problems in Penang Island, Malaysia (n = 256) (Na 2001)

1. Location of survey?	8. How do you get rid of the ants?
Urban 47%	Crush them with finger 5%
Suburban 41%	Aerosol spray 65%
Rural 12%	Hot water 2%
2. Age of respondents?	Sweep with a broom 10%
<20 6%	Ant bait 6%
20–29 20%	Not bothered at all 3%
30–39 20%	Other methods (eg., miracle chalk) .. 9%
40–49 29%	9. Best way to control ants?
50–59 16%	Aerosol spray 55%
60–69 6%	Sanitation 13%
70 and above 3%	Ant bait 8%
3. Gender of respondents?	Cooperation from neighbours 0%
Male 41%	Do not know 11%
Female 59%	Others (eg., miracle chalk) 13%
4. Problem with pest ants?	10. Main reason for ant control?
Yes 62%	Embarrassed in presence of guest ... 1%
No 38%	Ants transmit germs 4%
5. Do you find ants daily in your house?	Ants in food/drinks 38%
Yes 62%	Aesthetical nuisance 11%
No 38%	Others 27%
6. What causes ants in your house?	Not bothered 19%
Food/beverages 70%	11. Which is the most important?
Rubbish 5%	Ant problem 3%
other sources 14%	Rat problem 11%
do not know 11%	Leaking pipe 22%
7. Level of tolerance (# ants per sight)?	A crying baby 59%
1–10 30%	Unwashed clothes 5%
10–20 16%	12. Can ants be eliminated?
20–50 17%	Yes 29%
50–100 23%	No 65%
>100 14%	Not sure 6%

most likely reason. Sixty-three percent of the respondents had a tolerance of < 50 ants per sighting, while only 36 respondents out of 256 tolerated >100 ants per sighting. A low tolerance of homeowners to ants did not cause them to regard ants as a problem in their houses ($\chi^2 = 62.72$, $P < 0.05$, $df = 3$). However, if ants were found daily in the home, residents tolerated seeing more than 20 ants before initiating a control program ($\chi^2 = 7.51$, $P > 0.05$, $df = 3$).

Most homeowners (65%) used aerosol sprays for ant control; 6% used ant baits. Responses to another question, on the best way to control ants, indicated that only 55% chose aerosol sprays, indicating that many have experienced control failure using aerosols. Only 13% and 8%, respectively, of the respondents felt that sanitation and ant bait, respectively, were the best methods for ant control, while 55% commented that ant bait products were expensive when compared to aerosol. The majority (>80%) were not aware of the importance of sanitation in preventing ant problems in their houses.

Thirty-eight percent reported the main reason for ant control was the presence of ants in their food and beverages, while 11% initiated control when ants became an aesthetical nuisance. Only 4% of the respondents controlled ants because they were potential mechanical vectors of

pathogenic organisms. Surprisingly, 19% of the respondents were not bothered by the presence of ants. The importance of the ant problem in relation to several other negative situations was also investigated. Results indicated that ant problems received the lowest ranking when compared to other situations such as a crying baby, rat problems, leaking pipes, and unwashed clothes. Most homeowners (65%) felt that it would be impossible to eliminate pest ants from their premises.

Microorganisms From Ants in Food Outlets

Problems associated with ant infestations include food contamination (Lee et al., 2001b), stings, bites, and allergies (Goddard, 1993; Williams et al., 2001), contamination of surgical instruments in hospitals (Beatson, 1972), as well as serving as mechanical vectors of human diseases (Aleksev et al., 1972; Beatson, 1972; Edwards and Baker, 1981; Eichler, 1990; Bueno and Fowler, 1994). Most of the above studies concentrated on ants infesting hospital buildings, or on fire ants. There is limited information on microorganisms isolated from household ants found in food outlets, such as restaurants, cafeterias, and household kitchens. This study was initiated to provide information in this area.

Ants were collected on index cards (6.2 x 7.5 cm) and brought back to the laboratory in sterile petri dishes. In the laboratory, live ants were randomly selected and introduced into agar nutrient and Sabourand dextrose agar plates, and incubated at 37° C for 24-48 hours, and for 1-7 days, respectively. Colonies found growing on the agars were isolated and subcultured. Preliminary identification of bacteria was done according to the Gram's method. Gram negative bacteria were identified using *in vitro* API 20E and API 20NE diagnostic kits. Isolates of yeasts and fungi were stained with lactophenol cotton blue, and identified manually using a light microscope.

Table 6. Microorganisms isolated from pest ants collected in food-outlets and food preparation areas in Penang, Malaysia (Lim 2001)

Pathogen group and Species	Isolation frequency	Pathogen group and Species	Isolation frequency
Bacteria		Bacteria, continued	
Gram positive, bacilli	+++++	<i>Pseudomonas aeruginosa</i>	+
Gram positive, cocci	+++++	<i>Serratia liquefaciens</i>	+
Gram positive, streptococci	++	<i>Serratia plymuthica</i>	+
Gram positive, staphylococci	++	<i>Sphingomonas paucimobilis</i>	++
Gram negative, bacilli	++++	<i>Stenotrophomonas maltophilia</i>	++
Gram negative, cocci	+++	<i>Vibrio alginolyticus</i>	+
Gram negative, coccobacilli	+	<i>Vibrioparahaemolyticus</i>	++
<i>Acinetobacter baumannii</i>	+	<i>Vibrio fluvialis</i>	+
<i>Aeromonas hydrophila</i>	++++	<i>Yersinia pseudotuberculosis</i>	++
<i>Aeromonas salmonicida</i>	++	Fungi	
<i>Agrobacterium radiobacter</i>	++++	<i>Aspergillus flavus</i>	++++
<i>Alcaligenes faecalis</i>	+	<i>Aspergillus fumigatus</i>	++
<i>Bordetella bronchiseptica</i>	+	<i>Aspergillus niger</i>	++++
<i>Burkholderia cepacia</i>	++	<i>Cladosporium wernickii</i>	+
<i>Burkholderia pseudomallei</i>	+	<i>Fusarium oxysporum</i>	+
<i>Chryseobacterium meningosepticum</i>	++	<i>Penicillium</i> spp.	+
<i>Chryseomonas luteola</i>	+	<i>Scopulariopsis</i> spp.	++
<i>Erwinia</i> sp.	+	<i>Syncephalastrum</i> spp.	+
<i>Enterobacter sakazakii</i>	+	Yeast	
<i>Pantoea</i> spp.	+	<i>Geotrichum candidum</i>	+++
<i>Pasteurella multocida</i>	+	<i>Trichosporon cutaneum</i>	++

A total of 193 bacterial isolates were obtained; 55.4% of them were gram positive, and most of these were bacilli and cocci. Among the 23 species of bacteria identified, *Aeromonas hydrophila* and *Agrobacterium radiobacter* were the most frequently isolated (Table 6). Of the eight species of fungi isolated, *Aspergillus flavus* and *Aspergillus niger* were the most common. Two species of yeast were identified. This study provided an insight into various microorganisms found on the external body surface of household ants collected in food preparation premises.

Foraging Behavior

Food baits used on index cards were different in attractiveness towards common species of household ants. Field studies revealed that peanut butter was more attractive than honey to *M. pharaonis*, *M. destructor*, *Pheidole* sp., and *Solenopsis geminata*, while *T. melanocephalum* and *P. longicornis* preferred the latter (Table 7). Further studies conducted on a field population of *M. pharaonis* showed a seasonal preference to peanut butter and honey baits. In an 18-month study (February 1999 to August 2000), the attractancy to peanut butter and honey was found to vary. Granovsky and Howell (1983) also reported changes in bait preference. This behavior can be caused by changes in colony development (Erpenbeck, 1981), or food satiation (Edwards and Abraham, 1990). Regulation of nutrient intake is important to the foraging workers because queens and larvae have different nutritional requirements (Chong, 1997).

In the process of baiting household ants, it was regularly observed that different bait bases have varying attractiveness to foraging ants. The time for ants to become attracted to the bait can

Table 7. Food preference of several species of household ants as baited using ruled index cards

Species	Peanut butter*	Honey*
<i>Monomorium pharaonis</i>	++++	+
<i>Monomorium destructor</i>	++++	+
<i>Tapinomelanocephalum</i>	++	+++
<i>Pheidole</i> sp.	++++	+
<i>Paratrechina longicornis</i>	+	++++
<i>Solenopsis geminata</i>	++++	

*A '+' represents 20% of total ants feeding on the food attractant.

Table 8. Changes in food preference of a *Monomorium pharaonis* population in the field¹

Date	Peanut butter ²	Honey ²
Feb '99	+++++	
Jun '99	++++	+
Aug '99	+ +++++	
Dec '99	+++++	
Feb '00	++++	+
May '00	++++	+
Aug '00	+ +++++	

¹Experiment was conducted on one field population of *M. pharaonis* with three replicates (one replicate was done every alternate day).

²One '+' represents 20% of total ants feeding on the food attractant.

also vary with different formulations. Experiments were conducted using laboratory colonies of household ants. Depending on ant species and availability, all colonies have between 2-10 queens, 500-1000 workers, and 1-3 gm of brood. Blank bait bases containing 30% sucrose were formulated in the laboratory. It was previously determined that a 20-30% sucrose solution was the most attractive concentration for the majority of Malaysian household ant species (Lee, unpublished data). Choice experiments were done by presenting different bait bases simultaneously to foraging ants.

Results indicated that a liquid base was the most attractive base to all species tested. *Pheidole* spp. showed a relatively low response to all three bait bases, followed by gel base (Table 9). It was observed that granular toxicant bait gave better control against *Pheidole* sp. when compared to other toxicant bait bases (Lee, unpublished data). Paste base, which is widely used in commercial bait, received little or no response from the foraging ants. However, unlike the 30% sucrose used in this study, most commercial bait formulations contained one to several food attractants.

Studies were done to determine the speed of response of different species of household ants to liquid bait base. Using laboratory colonies in test arenas (40 x 25 cm), it was observed that *P. longicornis* registered the shortest time, 10 min, to achieve peak foraging activity, followed by *T. melanocephalum*, 20 min, when compared to other ant species (Table 10). The erratic and ran-

Table 9. Preference of laboratory colonies of several household ant species to different blank bait bases (at 30 minutes after introduction) (Beh, 2002)

Species	bait base		
	Liquid	Gel	Paste
<i>Monomorium pharaonis</i>	+++	+	-
<i>Monomorium floricola</i>	+++	+	+
<i>Monomorium destructor</i>	+++	+	-
<i>Paratrechina longicornis</i>	+++	++	-
<i>Pheidole</i> sp.	+	-	-
<i>Solenopsis molesta</i>	+++	++	-
<i>Formica</i> sp.	+++	+	+
<i>Tapinomamelanocephalum</i>	+++	++	-

A '+' represents 20% of total ants feeding on the bait formulation (or carrying the granular formulation).

Table 10. Differential response of laboratory colonies of several household ant species to liquid bait formulation (Beh, 2002)

Species	Time taken to achieve peak foraging activity (min)
<i>Monomorium pharaonis</i>	60
<i>Monomorium floricola</i>	100
<i>Monomorium destructor</i>	110
<i>Paratrechina longicornis</i>	10
<i>Solenopsis molesta</i>	80
<i>Formica</i> sp.	40
<i>Tapinomamelanocephalum</i>	20

dom foraging activity of both species could be a factor that increased the chance of the foraging ants finding the food source.

When screening residential premises to determine suitable houses to conduct the ant bait trials, inconsistent sampling results were often recorded when sampling time varied. Household ants may have a daily foraging activity pattern, especially those species that nest outdoors, but forage indoors for food. In this study, three ant species were chosen: *P. longicornis*, *M. pharaonis*, and *S. geminata*. These were infesting a large building, and samples were taken along the exposed outside corridor. Baited index cards were placed at locations where the ants were seen trailing every 2 hours and continuously up to 48 hours. The number of foraging workers on the index cards was counted at about 30 min after placement. Experiments were replicated three times and each replicate was done every alternate week. Results indicated that all three species have relatively similar foraging pattern (Figure 1). Foraging activity peaked 2-4 h after sunset (22:00), and the activity gradually ceased at about 15:00 in the afternoon. The findings suggested that foraging activity patterns of *M. pharaonis*, *P. longicornis*, and *S. geminata* were negatively correlated with ambient temperature. In Malaysia, the outdoor temperature averages 25°C at night, while day temperature averages 33°C, but can reach as high as 35°C. Hooper and Rust (1997) reported that foraging activity of *Solenopsis xyloni* began approximately 4 h before sunset and maximal activity occurred 2-7 h after sunset. They found that foraging workers avoid the times of the day when ground surface temperatures were relatively higher (Hooper and Rust, 1997).

Baiting Tropical Household Ants

Residual insecticide sprays are used for household ant control by Malaysian pest control operators (Chong et al., 1998; Lee et al., 1997). Because the number foraging accounts for only about 10% of total nest population (Adams et al., 1999), colony elimination using this method is not




Figure 1. Foraging activity patterns of *Paratrechina longicornis*, *Monomorium pharaonis*, and *Solenopsis geminata*

possible unless the nest is located and directly treated (Lee and Robinson, 2001). Residual spraying with pyrethroid insecticides has limitations in control, including unpredictable efficacy due to heterogeneity of treatment surface (Knight and Rust, 1990), insecticide repellency (Chong and Lee, 1999; Lee et al., 2002), and inability to eliminate the colony (Forschler and Evans, 1994a). Numerous bait toxicants evaluated have shown efficacy against several ant species such as the Argentine ant, odorous house ant, carpenter ant, Pharaoh ant, and fire ant. These toxicants are generally neurotoxic insecticides (Newton and Coombes, 1987; Oi et al., 1996; Lee, 2000), stomach poisons (Klotz et al., 1996, 1997a, 1997b, 2000a), metabolic inhibitors (Williams and Whelan, 1992; Forschler and Evans, 1994a, 1994b; Oi et al., 1994; Blachly and Forschler, 1996; Klotz et al., 2000b; Lee, 2000), and insect growth regulators (Edward and Clarke, 1978; Rupes et al., 1978; Newton, 1980; Oi et al., 1996, 2000; Reimer et al., 1991; Vail and Williams, 1995; Vail et al., 1996; Williams, 1990; Williams and Whelan, 1992; Williams and Vail, 1993, 1994; Williams et al., 1999).

Field studies were conducted to evaluate the performance of bait formulations against major household ant species: *M. pharaonis*, *T. melanocephalum*, *P. longicornis*, and *Pheidole* sp. All bait formulations containing fipronil and imidacloprid, hydramethylnon and boric acid, and sodium borate showed efficacy, with >75% reduction within a week post-treatment, with the exception of the borate-based bait against *Pheidole* sp. (Table 11). Bait containing methoprene eliminated the colonies by 8 weeks post-treatment. Horwood (1988) reported control of a field population of *Pheidole megacephala* using 0.5% methoprene in peanut butter.

Although a substantial reduction of household ants was recorded by most bait toxicants, limited success was registered when baiting crazy ants and ghost ants with paste and granular bait formulations. Hedges (1998) also reported that *P. longicornis* and *T. melanocephalum* were difficult to control with bait.

The effects of sanitation on field performance of toxic bait against household ants was also studied. Houses chosen for the trial were rated for their sanitary condition as follow: 1 = poor, 2 = moderate, and 3 = good. Two bait formulations were used: containerized bait and gel bait. Three containerized baits, each containing 1.65 g bait, were placed in houses, while 55 dabs of 0.09 g gel bait were applied in the other houses chosen for gel bait treatment. There was a positive correla-

Table 11. Summary of field studies on the performance of bait formulations against several species of Malaysian household ants

Bait active ingredient	Targetted species ¹	% reduction after n th week			
		1-week	4-week	8-week	12-week
hydramethylnon (1%) ²	Mp, Tm, Pl	82.0	83.7	54.9	-
fipronil (0.01%) ²	Mp, Tm, Pl	92.7	90.4	79.3	-
methoprene (0.5%) ³	Mp	51.9	87.6	100	100
hydramethylnon (1%) ⁴	Mp	89.1	89.9	100	100
boric acid (5.3%), sodium borate (4.3%) ⁴	Mp	76.2	74.5	78.2	-
boric acid (5.3%), sodium borate (4.3%) ⁵	Ph.	39.1	65.2	98.9	-
methoprene (0.5%) ⁶	Ph	6.5	94.3	100	100
imidacloprid (2.15%) ⁶	Mo	92.1	93.2	84.8	-
hydramethylnon (1%) ⁶	Mo	83.7	98.7	82.8	-

¹Mp = *Monomorium pharaonis*, Tm = *Tapinoma melanocephalum*, Pl = *Paratrechina longicornis*, Ph = *Pheidole* sp., Mo = *Monomorium* spp. (*destructor*, *floricola*, and *pharaonis*).

²Based on Lee (2000).

³Based on Lee et al. (2001).

⁴Based on Lee and Lee (2001).

⁵Based on Loke (2002).

⁶Lee, unpublished data.




Figure 2. Relationship between containerized bait performance and sanitary condition in baited premises.




Figure 3. Relationship between gel bait performance and sanitary condition in baited premises.

tion ($y = 49.2 \pm 16.9x$; $x \geq 1$, $r^2 = 0.849$, $P = 0.0038$) between sanitation and the performance of containerized bait (Figure 2). However, no correlation ($r^2 = 0.2659$, $P = 0.4589$) was found between sanitary condition and the performance of gel bait (Figure 3). This result suggests that sanitation did not play a significant role when baiting ants using gel bait, perhaps because of the higher number of placements in gel treatment when compared to containerized bait that had only three placements. Therefore, by increasing bait placements, it may be possible to overcome efficacy problems due to bad sanitary conditions. A similar finding was also observed earlier when baiting American cockroaches in urban slums (Lee, unpublished data).

In the tropics several ant species of ants may occur within a living unit at the same time. One species may be predominant and be more aggressive, and have a wider foraging territory than others. One bait may show good performance against one species, but may not be effective against other household ant species. When baiting *Monomorium pharaonis*, *M. destructor*, and *M. floridicola* using imidacloprid gel bait, a significant reduction (>90%) in the numbers of *Monomorium* spp. was recorded at 1-2 weeks post-treatment (Figure 4). However, following a reduction of *Monomorium* spp., an increase in *T. melanocephalum* and *P. longicornis* counts occurred on index cards. It is speculated that upon reduction or elimination of the more dominant *Monomorium* spp., other tramp ant species, such as *T. melanocephalum* and *P. longicornis*, were now able to forage at a wider area for food and thus were found in larger numbers on index cards. What long term effects the imidacloprid gel bait has on these species is still unknown. However, it is likely that the effect would be minimal, because it was observed that the two species were not receptive to the freshly applied gel bait.

SUMMARY and CONCLUSION

Ants are an increasingly important group of household pests in the human environment in Malaysia. This was evident from the various household pest surveys that had been conducted on




Figure 4. Changes in number of foraging ants upon baiting *Monomorium* spp. with 2.15% imidacloprid gel bait.

homeowners in Penang, Malaysia. Twenty-five species were found in and around buildings and structures. Most of these were tramp ants, such as *Monomorium* spp., *P. longicornis*, *T. melanocephalum*, and *Pheidole* sp. Most tropical household ants prefer a liquid bait base over gel and paste base. However, a liquid bait is susceptible to spillage and desiccation. Despite a liquid bait being the preferred base choice, all granular, paste, and gel baits tested showed efficacy against field colonies of household ants. Most formulations showed >70% reduction in ant numbers after 4 weeks post-baiting. It was also noted that the effect of sanitation on bait performance in the premises baited can be reduced if a higher bait placement was done. More studies, especially on foraging biology of important tropical household ants such as *M. destructor*, *Pheidole* sp., *T. melanocephalum*, and *P. longicornis*, are warranted, so that a reliable pest management program can be effectively constructed.

ACKNOWLEDGMENTS

I thank William H. Robinson and N.L. Chong for comments and proof reading the draft manuscript; I. Ridzuan, H.H. Yap, J. Ho, and T. Fernandez for information on household insecticide products and the pest control industry; I. Darah and K.K. Peh for helping with microorganism identification and bait formulation; L.C. Lee, A.G.H. Eow, J.P.S. Na, M.Y. Low, S.W. Beh, P.Y. Loke, K.T. Koay, K.T. Lim, S.H. Looi, P.I. Ong, C.C. Tan, G.K. Tan, Y. Way, F.L. Yap, K.L. Yee, and L.H. Yee for their work on various projects; and Bayer AG Germany, Aventis Environmental Health (Malaysia), Dow AgroSciences (USA), Clorox Technical Center (USA), and Reckitt Benckiser (Australia) for research grants on household ants.

REFERENCES

- Adams, A., Kunkel, S., Dodd, G., and Hobel, S. 1999. Method and procedure for evaluating biological performance of Pharaoh ant, *Monomorium pharaonis* (Hymenoptera: Formicidae) baits. In: Robinson, W.H., Rambo, G. and Rettich, F., eds. Proceedings of the Third International Conference on Urban Pests. Prague: Graficke zavody Hronov., pp. 203-209.
- Aleksev, A.N., Bibivoka, V.A., Brikman, T., and Kantarbaeva, Z. 1972. The persistence of viable plague microbes on the epidermis and in the alimentary tract of *Monomorium pharaonis* in experimental conditions. Med. Parazitol. Paraz. Bole. 41: 237-239.
- Beaton, S.H. 1972. Pharaoh's ants as pathogen vectors in hospitals. Lancet 1: 425-427.
- Beh, S.W. 2002. Studies on bait matrices and formulations for development of ant baits against eight species of household ants (Hymenoptera: Formicidae). B. Appl. Sci. Dissertation. Penang: School of Biological Sciences, Universiti Sains Malaysia. 66 pp.
- Blachly, J.S. and Forschler, B.T. 1996. Suppression of late-season Argentine ant (Hymenoptera: Formicidae) field populations using a perimeter treatment with containerized baits. J. Econ. Entomol. 89: 1497-1500.
- Bueno, O.C. and Fowler, H.G. 1994. Exotic ants and native ant fauna of Brazilian hospitals. In: Williams, D.F., ed. Exotic ants – Biology, impact and control of introduced species. Boulder: Westview Press, pp. 191-198.
- Chong, A.S.C. 1997. Feeding behaviour of the Pharaoh ant, *Monomorium pharaonis* (L.) (Hymenoptera: Formicidae): Recruitment, food selection and distribution. M.Sc. thesis. Penang: School of Biological Sciences, Universiti Sains Malaysia. 129 pp.
- Chong, A.S.C. and Lee, C.Y. 1999. Household ants. In: Lee, C.Y., Yap, H.H., Chong, N.L. and Jaal, Z., eds. Urban Pest Control – A Malaysian Perspective. Penang: Universiti Sains Malaysia Press, pp. 66-80.
- Chong, A.S.C., Gabriel, L.C., Lee, C.Y., and Yap, H.H. 1998. Contact toxicities of etofenprox and bendiocarb against the odorous house ant (*Tapinoma sessile* Say). Malay. Appl. Biol. 27: 127-129.
- Edwards, J.P. 1986. The biology, economic importance, and control of the Pharaoh's ant, *Monomorium pharaonis* (L.). Pp. 257-271 in: Vinson, S.B., ed. Economic impact and control of social insects. New York: Praeger.
- Edwards, J.P. and Abraham, L. 1990. Changes in food selection by workers of the pharaoh's ant, *Monomorium pharaonis*. Med. Vet. Entomol. 4: 205-211.
- Edwards, J.P. and Baker, L.F. 1981. Distribution and importance of the Pharaoh's ant, *Monomorium pharaonis* L., in national health service hospitals in England. J. Hosp. Infec. 2: 249-254.
- Edwards, J.P. and Clarke, B. 1978. Eradication of Pharaoh's ants with baits containing the insect juvenile

- hormone analogue methoprene. *Int. Pest Control* 17: 5-10.
- Eichler, W. 1990. Health aspects and control of *Monomorium pharaonis*. In: Vander Meer, R.K., Jaffe, K. and Cedeno-Leon, A., eds. *Applied Myrmecology: A World Perspective*. Boulder: Westview Press, pp. 671-675.
- Erpenbeck, A. 1981. Nahrungseintrag und Verteilung in Sozietaten der Pharaoameise *Monomorium pharaonis* (L.) (Hymenoptera: Formicidae) in Abhängigkeit vom Brutzyklus. *Mutteilungen des Deutschen Gesellschaft für Angewandte Entomologie* 3: 87-89.
- Forschler, B.T. and Evans, G.M. 1994a. Argentine ant (Hymenoptera: Formicidae) foraging activity response to selected containerized baits. *J. Entomol. Sci.* 29: 209-214.
- Forschler, B.T. and Evans, G.M. 1994b. Perimeter treatment strategy against containerized baits to manage Argentine ants, *Linepithema humile* (Mayr) (Hymenoptera: Formicidae). *J. Entomol. Sci.* 29: 264-267.
- Hansen, L.D. and Akre, R.D. 1993. Urban pest management of carpenter ants. In: Wildey, K.B. and Robinson, Wm.H., eds. *Proceedings of the First International Conference on Insect Pests in the Urban Environment*. Exeter: BPCC Wheatons Ltd., pp. 271-279.
- Hedges, S.A. 1998. *Field guide for the management of structure-infesting ants*. Cleveland: G.I.E. Inc., Publishers. 304 pp.
- Holldobler, B. and Wilson, E.O. 1990. *The ants*. Cambridge: Harvard University Press. 732 pp.
- Hooper, L.M. and Rust, M.K. 1997. Food preference and patterns of foraging activity of the Southern fire ant (Hymenoptera: Formicidae). *Ann. Entomol. Soc. Am.* 90: 246-253.
- Horwood, M.A. 1988. Control of *Pheidole megacephala* (F.) (Hymenoptera: Formicidae) using methoprene baits. *J. Aust. Entomol. Soc.* 27: 257-258.
- Goddard, J. 1993. *Physician's Guide to Arthropods of Medical Importance*. Atlanta: CRC Press. 332 pp.
- Gooch, H. 1999. Baiting remains the treatment of choice. *Pest Control* 67 (9): 40-43.
- Granovsky, T.A. and Howell, H.H. 1983. Texas A&M research team develops new Pharaoh's ant control technique. *Pest Control Technol.* 11: 30-34.
- Jenkins, M. 2001. Battling a common enemy. *Pest Control* 69 (9): S10-S12, S16, S27.
- Kaminski, L. 2000. Ants are still tops, for now. *Pest Control* 68 (9): S9-S11.
- Klotz, J.H., Oi, D.H., Vail, K.M., and Williams, D.F. 1996. Laboratory evaluation of a boric acid liquid bait on colonies of *Tapinoma melanocephalum*, Argentine ants and Pharaoh ants (Hymenoptera: Formicidae). *J. Econ. Entomol.* 89: 673-677.
- Klotz, J.H., Vail, K.M., and Williams, D.F. 1997a. Toxicity of a boric acid-sucrose water bait to *Solenopsis invicta* (Hymenoptera: Formicidae). *J. Econ. Entomol.* 90: 488-491.
- Klotz, J.H., Vail, K.M., and Williams, D.F. 1997b. Liquid boric acid bait for control of structural infestations of Pharaoh ants (Hymenoptera: Formicidae). *J. Econ. Entomol.* 90: 523-526.
- Klotz, J.H., Greenberg, L., Amrhein, C., and Rust, M.K. 2000a. Toxicity and repellency of borate-sucrose water baits to Argentine ants (Hymenoptera: Formicidae). *J. Econ. Entomol.* 93: 1256-1258.
- Klotz, J.H., Greenberg, L., and Venn, G. 2000b. Evaluation of two hydramethylnon granular baits for control of Argentine ant (Hymenoptera: Formicidae). *Sociobiol.* 36: 201-207.
- Knight, R.L. and Rust, M.K. 1990. Repellency and efficacy of insecticides against foraging workers in laboratory colonies of Argentine ants (Hymenoptera: Formicidae). *J. Econ. Entomol.* 83: 1402-1408.
- Lee, C.Y. 2000. Performance of hydramethylnon- and fipronil-based containerized baits against household ants in residential premises. *Trop. Biomed.* 17: 45-48.
- Lee, C.Y. and Lee, L.C. 2001. Field and laboratory evaluation of a boron-based containerised dual-bait formulation against the Pharaoh ant, *Monomorium pharaonis* (L.) (Hymenoptera: Formicidae). *Trop. Biomed.*: In press.
- Lee, C.Y. and Robinson, W.H. 2001. *Handbook of Malaysian Household and Structural Pests*. Kuala Lumpur: Pest Control Association of Malaysia. 96 pp.
- Lee, C.Y., Eow, A.G.H., Na, J.P.S., Lee, L.C., Lee, K.M., Ngee, P.S., and Lim, S.Y. 2002. Managing pest ants in urban environment. *PCAM Pest Info*: In press.
- Lee, C.Y., Lee, L.C., Na, J.P.S., Loke, P.Y., Lim, K.T., and Teo, E.H.H. 2001b. Field performance of a methoprene granular bait against the Pharaoh ant, *Monomorium pharaonis* (Hymenoptera: Formicidae). *J. Biosci.*: In press.
- Lee, C.Y., Lim, C.Y., and Yap, H.H. 1997. Contact toxicity of bendiocarb against three species of Malaysian household ants (Hymenoptera: Formicidae). *J. Biosci.* 7: 79-82.
- Lee, C.Y., Lim, C.Y., and I. Darah. 2001a. Survey on structure-infesting ants (Hymenoptera: Formicidae) in food preparative outlets. *Trop. Biomed.*: In press.
- Lee, C.Y., Yap, H.H., Chong, N.L., and Jaal, Z. 1999. *Urban Pest Control – A Malaysian Perspective*. Penang: Universiti Sains Malaysia Press. 134 pp.

- Lim, C.Y. 2001. Isolation and identification of pathogenic microorganisms from household ants (Hymenoptera: Formicidae). B.Appl. Sci. dissertation. Penang: School of Biological Sciences, Universiti Sains Malaysia. 98 pp.
- Loke, P.Y. 2002. Foraging behavior of *Pheidole megacephala* (Fabricius) (Hymenoptera: Formicidae). B.Appl. Sci. dissertation. Penang: School of Biological Sciences, Universiti Sains Malaysia. 104 pp.
- Na, J.P.S. 2001. Studies on status, diversity, foraging behaviour and temperature/ humidity tolerance of household ant, *Tapinoma* spp. (Hymenoptera: Formicidae). B.Appl. Sci. dissertation. Penang: School of Biological Sciences, Universiti Sains Malaysia. 105 pp.
- Na, J.P.S. and Lee, C.Y. 2001. Identification key to common urban pest ants in Malaysia. Trop. Biomed.: In press.
- Newton, J. 1980. Alternatives to chlordecone for Pharaoh's ant control. Int. Pest Control 22: 112-114, 135.
- Newton, J. and Coombes, D.S. 1987. A comparison of a range of novel and conventional insecticides for Pharaoh's ant control. Int. Pest Control 29: 45-47.
- Oi, D.H., Vail, K.M., Williams, D.F., and Bieman, D.N. 1994. Indoor and outdoor foraging locations of Pharaoh ants (Hymenoptera: Formicidae) and control strategies using bait stations. Fla. Entomol. 77: 85-91.
- Oi, D.H., Vail, K.M., and Williams, D.F. 1996. Field evaluation of perimeter treatments for Pharaoh ant (Hymenoptera: Formicidae) control. Fla. Entomol. 79: 252-263.
- Oi, D.H., Vail, K.M., and Williams, D.F. 2000. Bait distribution among multiple colonies of Pharaoh ants (Hymenoptera: Formicidae). J. Econ. Entomol. 93: 1247-1255.
- Passera, L. 1994. Characteristics of tramp species. In: Williams, D.F., ed. Exotic ants – Biology, impact and control of introduced species. Boulder: Westview Press, pp. 23-43.
- Pisarki, B. 1982. Ants (Hymenoptera: Formicoidea) of Warsaw and Mazovia. Memor. Zool. 36: 73-90.
- Prins, A.J., Robertson, H.G., and Prins, A. 1990. Pest ants in urban and agricultural areas of Southern Africa. In: Vander Meer, R.K., Jaffe, K. and Cedeno-Leon, A., eds. Applied Myrmecology: A World Perspective. Boulder: Westview Press, pp. 25-33.
- Reimer, N.J., Beardsley, J.W., and Jahn, G. 1990. Pest ants in the Hawaiian islands. In: Vander Meer, R.K., Jaffe, K. and Cedeno, A., eds. Applied Myrmecology – A World Perspective. California: Westview Press, pp. 40-50.
- Reimer, N.J., Glancey, B.M., and Beardsley, J.W. 1991. Development of *Pheidole megacephala* (Hymenoptera: Formicidae) colonies following ingestion of fenoxycarb and pyriproxyfen. J. Econ. Entomol. 84: 56-60.
- Robinson, W.H. 1996. Urban Entomology – Insect and mite pests in the human environment. New York: Chapman & Hall. 430 pp.
- Rupes, V., Hrdy, I., Pinterova, J., Zdarek, J., and Kreck, J. 1978. The influence of methoprene on pharaoh's ant, *Monomorium pharaonis* colonies. Acta Entomol. Bohem. 75: 155-163.
- Taber, S.W. 2000. Fire ants. College Station: Texas A & M University Press. 308 pp.
- Thompson, C.R. 1990. Ants that have pest status in the United States. In: Vander Meer, R.K., Jaffe, K. and Cedeno, A., eds. Applied Myrmecology: A World Perspective. Boulder: California: Westview Press, pp. 51-67.
- Vail, K.M. and Williams, D.F. 1995. Pharaoh ant (Hymenoptera: Formicidae) colony development after consumption of pyriproxyfen baits. J. Econ. Entomol. 88: 1695-1702.
- Vail, K.M., Williams, D.F., and Oi, D.H. 1996. Perimeter treatment with two bait formulations of pyriproxyfen for control of Pharaoh ants (Hymenoptera: Formicidae). J. Econ. Entomol. 89: 1501-1507.
- Vega, S.J. and Rust, M.K. 2001. The Argentine ant – a significant invasive species in agricultural, urban and natural environments. Sociobiol. 37: 3-25.
- Wetterer, J.K., Miller, S.E., Wheeler, D.E., Olson, C.A., Polhemus, D.A., Pitts, M., Ashton, W., Himler, A.G., Yospin, M.M., Helms, K.R., Harken, E.L., Gallaher, J., Dunning, C.E., Nelson, M., Litsinger, J., Southern, A., and Burgess, T.L. 1999. Ecological dominance by *Paratrechina longicornis* (Hymenoptera: Formicidae), an invasive tramp ant, in Biosphere 2. Fla. Entomol. 82: 381-388.
- Williams, D.F. 1990. Effects of fenoxycarb baits on laboratory colonies of the Pharaoh's ant, *Monomorium pharaonis*. In: Vander Meer, R.K., Jaffe, K. and Cedeno-Leon, A., eds. Applied Myrmecology: A World Perspective. Boulder: Westview Press, pp. 676-683.
- Williams, D.F. and Whelan, P.M. 1992. Bait attraction of the introduced pest ant, *Wasmannia auropunctata* (Hymenoptera: Formicidae) in the Galapagos Islands. J. Entomol. Sci. 27: 29-34.
- Williams, D.F. and Vail, K.M. 1993. Pharaoh ant (Hymenoptera: Formicidae): fenoxycarb baits affect colony development. J. Econ. Entomol. 86: 1136-1143.
- Williams, D.F. and Vail, K.M. 1994. Control of a natural infestation of the Pharaoh ant (Hymenoptera: Formicidae) with a con grit bait of fenoxycarb. J. Econ. Entomol. 87: 108-115.

- Williams, D.F., Brenner, R.J., and Milne, D. 1999. Precision targeting: reduced pesticide use strategy for Pharaoh's ant (Hymenoptera: Formicidae) control. Pp. 195-201 in: Robinson, W.H., Rambo, G. and Rettich, F., eds. Proceedings of the Third International Conference on Urban Pests. Prague: Graficke zavody Hronov.
- Williams, D.F., Collins, H.L., and Oi, D.H. 2001. The red imported fire ant (Hymenoptera: Formicidae): An historical perspective of treatment programs and the development of chemical baits for control. *Am. Entomol.* 47: 146-159.
- Wood, F.E., Robinson, W.H., Kraft, S.K., and Zungoli, P.A. 1981. Survey of attitudes and knowledge of public housing residents toward cockroaches. *Bull. Entomol. Soc. Am.* 27: 9-13.
- Yap, H.H. 1988. Household pests and household insecticide usage in Malaysia. *MACA Newsletter* 1: 11-12.
- Yap, H.H. 1999. Insecticide usage in the urban environment. In: Lee, C.Y., Yap, H.H., Chong, N.L. and Jaal, Z., eds. *Urban Pest Control – A Malaysian Perspective*. Penang: Universiti Sains Malaysia Press, pp. 8-4.
- Yap, H.H. and Foo, A.E.S. 1984. Household pests and household insecticide usage on Penang Island, Malaysia – A questionnaire survey. *Bull. Pub. Hlth. Soc.* 16: 2-8.
- Yap, H.H. and Lee, C.Y. 1994. A preliminary study on the species composition of household ants on Penang Island, Malaysia. *J. Biosci.* 5: 64-66.
- Yap, H.H. and Lee, C.Y. 1996. Current status of pest control industry in Malaysia. *Malaysian Pest Control Seminar – Urban Pest Control for a Healthier Environment*. Pest Control Association of Malaysia. Genting Highlands, Malaysia. 22-23 March 1996.
- Yap, H.H., Lee, Y.W., Ong, C.H., Ridzuan, I., Quah, E.S., and Chong, N.L. 1999. The abundance and control of household pests in Penang, Malaysia – questionnaire and trapping survey. *J. Biosci.* 10: 62-69.